

MATA37 Final Review

$$\cdot \int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x \text{ where } \Delta x = \frac{b-a}{n} \quad x_i^* = a + (\Delta x)i$$

• The Darboux Definition

Let $a, b \in \mathbb{R}$, $a < b$. Suppose f is bounded on $[a, b]$, let $p = \{x_i\}_{i=0}^n$ be any partition of $[a, b]$

Then for each $i = 1, \dots, n$

$$m_i = \inf \{f(x) \mid x \in [x_{i-1}, x_i]\}$$

$$M_i = \sup \{f(x) \mid x \in [x_{i-1}, x_i]\}$$

$$U(f, p) = \sum_{i=1}^n M_i (x_i - x_{i-1})$$

$$L(f, p) = \sum_{i=1}^n m_i (x_i - x_{i-1})$$

$$\cdot f \text{ is integrable on } [a, b] \Leftrightarrow \sup \{L(f, p) \mid p \text{ is any partition of } [a, b]\} \\ = \inf \{U(f, p) \mid p \text{ is any partition of } [a, b]\}$$

• Integrability Reformulation

$$f \text{ is integrable on } [a, b] \Leftrightarrow \forall \epsilon > 0, \exists p \text{ partition of } [a, b] \text{ s.t. } U(f, p) - L(f, p) < \epsilon$$

After midterm

• Determine if an improper integral conv or div

① By definition Let $A \rightarrow \infty$ then $\lim_{A \rightarrow \infty} \int_a^A f(x) dx$

② Comparison Theorem (CT)

• If $0 \leq g(x) \leq f(x) \forall x \in [a, \infty)$ and $\int_a^\infty f(x) dx$ conv, then $\int_a^\infty g(x) dx$ conv

• If $0 \leq h(x) \leq g(x) \forall x \in [a, \infty)$ and $\int_a^\infty h(x) dx$ div, then $\int_a^\infty g(x) dx$ div

Proof: Given f, g are cont on $[a, \infty)$

Suppose $0 \leq g(x) \leq f(x) \forall x \in [a, \infty)$ ①

$\int_a^\infty f(x) dx$ converges ②

WTS $\int_a^\infty g(x) dx$ converges

$\Downarrow$
WTS $\lim_{A \rightarrow \infty} \int_a^A g(x) dx$ exists

Let $A \in [a, \infty)$ be arbitrary

① $\Rightarrow 0 \leq g(x) \leq f(x) \forall x \in [a, A]$

$\Rightarrow \int_a^A 0 dx \leq \int_a^A g(x) dx \leq \int_a^A f(x) dx$, by int properties

$\Leftrightarrow 0 \leq \int_a^A g(x) dx \leq \int_a^A f(x) dx \forall A \geq a$

$\Rightarrow \lim_{A \rightarrow \infty} 0 \leq \lim_{A \rightarrow \infty} \int_a^A g(x) dx \leq \lim_{A \rightarrow \infty} \int_a^A f(x) dx$

$\Leftrightarrow 0 \leq \lim_{A \rightarrow \infty} \int_a^A g(x) dx \leq \int_a^\infty f(x) dx$

$\underbrace{\lim_{A \rightarrow \infty} \int_a^A g(x) dx}_{\text{is an area accumulation func. Note it's increasing as } A \rightarrow \infty} \leq \underbrace{\int_a^\infty f(x) dx}_{\text{conv by ②} = \#}$

$\therefore$ This limit exists.

$\therefore \int_a^\infty g(x) dx$ conv

Given h, g are cont on $[a, \infty)$

Suppose $0 \leq h(x) \leq g(x) \forall x \in [a, \infty)$ ①

$\int_a^\infty g(x) dx$ diverges ②

WTS $\int_a^\infty g(x) dx$ diverges

$\Downarrow$
WTS $\lim_{A \rightarrow \infty} \int_a^A g(x) dx$ DNE

Let $A \in [a, \infty)$ be arbitrary

① $\Rightarrow 0 \leq h(x) \leq g(x) \forall x \in [a, A]$

$\Rightarrow \int_a^A 0 dx \leq \int_a^A h(x) dx \leq \int_a^A g(x) dx$, by int properties

$\Leftrightarrow 0 \leq \int_a^A h(x) dx \leq \int_a^A g(x) dx \forall A \geq a$

$\Rightarrow \lim_{A \rightarrow \infty} 0 \leq \lim_{A \rightarrow \infty} \int_a^A h(x) dx \leq \lim_{A \rightarrow \infty} \int_a^A g(x) dx$

$\Leftrightarrow 0 \leq \lim_{A \rightarrow \infty} \int_a^A h(x) dx \leq \int_a^\infty g(x) dx$

$\underbrace{\lim_{A \rightarrow \infty} \int_a^A h(x) dx}_{\text{div by ②} = \text{DNE}} \leq \underbrace{\int_a^\infty g(x) dx}_{\text{is an area accumulation func. Note it's increasing as } A \rightarrow \infty}$

$\therefore$ This limit DNE

$\therefore \int_a^\infty g(x) dx$ div

if $0 < p \leq 1$, then $\frac{1}{x^p} \geq \frac{1}{x}$ for $x \in [1, \infty)$ and $\int_1^\infty \frac{1}{x^p} dx$ div
 if $p > 1$, then $\frac{1}{x^p} < \frac{1}{x}$ for $x \in [1, \infty)$ and $\int_1^\infty \frac{1}{x^p} dx$ conv to $\frac{1}{p-1}$ } $[1, \infty]$
 if $0 < p < 1$, then $\frac{1}{x^p} < \frac{1}{x}$ for $x \in [0, 1]$ and $\int_0^1 \frac{1}{x^p} dx$ conv to $\frac{1}{1-p}$
 if $p \geq 1$, then $\frac{1}{x^p} \geq \frac{1}{x}$ for $x \in [0, 1]$ and $\int_0^1 \frac{1}{x^p} dx$ div } $[0, 1]$

e.g.

(i) Does $\int_3^\infty \left(\frac{\cos^2(x)+1}{\sqrt{1+x^6}} \right) dx$ conv or div?

① for $x \in [3, \infty)$, $g(x) = \frac{\cos^2(x)+1}{\sqrt{1+x^6}} > 0$

$$\int_3^\infty \frac{\cos^2(x)+1}{\sqrt{1+x^6}} dx \sim \int_3^\infty \frac{2}{\sqrt{x^6}} dx = \int_3^\infty \frac{2}{x^3} dx \text{ conv}$$

Consider $\int_3^\infty f(x) dx$

$$= \lim_{A \rightarrow \infty} \int_3^A \frac{2}{x^3} dx$$

$$= \lim_{A \rightarrow \infty} -x^{-2} \Big|_3^A$$

$$= \lim_{A \rightarrow \infty} -\left(\frac{1}{A^2} - \frac{1}{9}\right)$$

$$= \frac{1}{9} \quad \therefore \text{lim exists}$$

$\therefore \int_3^\infty f(x) dx$ conv

$\therefore$ By CT, $\int_3^\infty \frac{\cos^2(x)+1}{\sqrt{1+x^6}} dx$ conv.

② Find a good & explicit comparison

- bigger
- easier
- conv

$$\text{for } x \in [3, \infty), g(x) = \frac{\cos^2(x)+1}{\sqrt{1+x^6}}$$

$$\leq \frac{1+1}{\sqrt{1+x^6}} \quad \text{max numerator b/c } \cos^2(x) \leq 1 \forall x$$

$$= \frac{2}{\sqrt{1+x^6}} \leq \frac{2}{\sqrt{x^6}} \quad \text{min denominator}$$

$$= \frac{2}{x^3} = f(x)$$

(ii) Does $\int_0^1 \frac{\csc^2(x)}{x^{\frac{3}{2}}} dx$ conv or div

① For $x \in [0, 1]$, $g(x) = \frac{1}{\sin^2(x)x^{\frac{3}{2}}} \geq 0$

② Find a good & explicit comparison

$$Rv \sim \int_0^1 \frac{1}{x^{\frac{3}{2}}} dx$$

- smaller
- easier
- div

$$\text{For } x \in [0, 1], g(x) = \frac{1}{\sin^2(x)x^{\frac{3}{2}}} \geq \frac{1}{1 \cdot \frac{3}{2}} \quad \text{max denominator b/c } 0 < \sin(x) \leq 1 \forall x$$

$$\left(x^{-\frac{3}{2}} \right) h(x)$$

$$= \int_0^1 \frac{1}{\sin^2(x)x^{\frac{3}{2}}} dx$$

$$\text{consider } \int_0^1 h(x) dx = \int_0^1 x^{-\frac{3}{2}} dx$$

$$= \lim_{A \rightarrow 0^+} \int_A^1 x^{-\frac{3}{2}} dx$$

$$= \lim_{A \rightarrow 0^+} \left. \frac{x^{-\frac{1}{2}}}{-\frac{1}{2}} \right|_A^1$$

$$= \lim_{A \rightarrow 0^+} -2 \left(1 - \frac{1}{\sqrt{A}} \right)$$

$$= +\infty \quad \therefore \text{lim does not exist}$$

$\therefore \int_0^1 h(x) dx$ div

$\therefore \int_0^1 \frac{\csc^2(x)}{x^{\frac{3}{2}}} dx$ div by CT.

Definition: Given $\{a_n\}$ let $L \in \mathbb{R}$. We say $\{a_n\}$ converges to L

iff $\exists L \in \mathbb{R}, \forall \epsilon > 0, \exists N > 0$ ^{a huge number} s.t. for all $n \in \mathbb{N}$
if $n > N$ then $|a_n - L| < \epsilon$

Prove a sequence is conv we have two ways

① ϵ - N proof

② Bounded Monotone Convergent Theorem (BMCT)

e.g. of ①

(i) Prove $a_n = \frac{n^2 - 2}{n^2 + 2n + 2}$ converges

wts $\exists L \in \mathbb{R}, \forall \epsilon > 0, \exists N > 0$ s.t. all for $n \in \mathbb{N}$, if $n > N$ then $|a_n - L| < \epsilon$

Pf

choose $L = 1 \in \mathbb{R}$ (compute $\lim_{n \rightarrow \infty} a_n$)

Let $\epsilon > 0$ be arbitrary

choose $N = \frac{2}{\epsilon} > 0$

Suppose $n > N$

$$\text{then } \left| \frac{n^2 - 2}{n^2 + 2n + 2} - 1 \right| = \left| \frac{n^2 - 2 - n^2 - 2n - 2}{n^2 + 2n + 2} \right|$$

$$= \left| \frac{-2n - 4}{n^2 + 2n + 2} \right| = \frac{2 \cdot |n + 2|}{|n^2 + 2n + 2|} = \frac{2 \cdot |n + 2|}{|n^2 + 2n + 2|} = \frac{2(n + 2)}{n^2 + 2n + 2} \text{, by alg. abs value property}$$

$$\text{we have } |a_n - L| = \frac{2(n + 2)}{n^2 + 2n + 2} < \frac{2(n + 2)}{n^2 + 2n} \text{ (min denom)} = \frac{2(n + 2)}{n(n + 2)} = \frac{2}{n} < \frac{2}{N} = \frac{2}{\frac{2}{\epsilon}} = \epsilon \text{ as needed}$$

WANT: $|a_n - L| < \frac{\epsilon}{2}$
PER, $\epsilon > 0$

given: $n > N$

$$\Rightarrow \frac{1}{n} < \frac{1}{N}$$

$$\Rightarrow \frac{2}{n} < \frac{2}{N}$$

$$\frac{2}{N} = \epsilon$$

$$2 \cdot \frac{1}{N} = \epsilon$$

$$\frac{1}{N} = \frac{\epsilon}{2}$$

$$N = \frac{2}{\epsilon}$$

(ii)

Ex2 Prove that $\underbrace{\left\{ \frac{2n-1}{n-3} \right\}}_{a_n}$ conv to 2

Wts $\forall \epsilon > 0, \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$ then $\left| \frac{2n-1}{n-3} - 2 \right| < \epsilon$

Pf let $\epsilon > 0$ be arbitrary

choose $N = \max\{4, \frac{20}{\epsilon}\} > 0$

Suppose $n > N$

$$\text{Then } \left| \frac{2n-1}{n-3} - 2 \right| = \left| \frac{2n-1-2n+6}{n-3} \right| = \left| \frac{5}{n-3} \right| = \frac{5}{|n-3|} = 5 \cdot \frac{1}{|n-3|} = 5 \cdot \frac{1}{n(1-\frac{3}{n})} = 5 \cdot \frac{1}{n} \cdot \frac{1}{|1-\frac{3}{n}|} \quad n=0, n=3 \text{ are problem pts}$$

★ if $n > 4$ (★ $N \geq 4$)

$$\Rightarrow \frac{1}{n} < \frac{1}{4}$$

$$\Rightarrow \frac{3}{n} < \frac{3}{4}$$

$$\Rightarrow 1 - \frac{3}{n} > 1 - \frac{3}{4} = \frac{1}{4} \quad \therefore \left| 1 - \frac{3}{n} \right| > \frac{1}{4} \Rightarrow \frac{1}{|1-\frac{3}{n}|} < 4$$

$$\text{Therefore } \left| \frac{2n-1}{n-3} - 2 \right| = 5 \cdot \frac{1}{n} \cdot \underbrace{\frac{1}{|1-\frac{3}{n}|}}_{< 4} < 5 \cdot \frac{1}{n} \cdot 4 = \frac{20}{n} < \frac{20}{N} < \frac{20}{(\frac{20}{\epsilon})} = \epsilon \quad \text{as needed}$$

(iii) Prove $\{1 + (-1)^n\}$ diverges

Assume $\{1 + (-1)^n\}$ converges to some $L \in \mathbb{R}$

We have $\forall \epsilon > 0, \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$ then $|a_n - L| < \epsilon$

Take $\epsilon = 1 > 0$

We have $\exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$, then $|1 + (-1)^n - L| < 1$

Case 1 n is odd: if $n > N \Rightarrow |0 - L| < 1 \Rightarrow |L| < 1$
 $\Leftrightarrow -1 < L < 1$

Case 2 n is even: if $n > N \Rightarrow |2 - L| < 1 \Rightarrow 1 < 2 - L < 1$

$$\Leftrightarrow -3 < -L < -1$$

$$\Leftrightarrow 1 < L < 3$$

$$\text{so } (L \in (-1, 1) \cap (1, 3)) = \emptyset$$

contradiction

$\therefore \{1 + (-1)^n\}$ diverges

(iv) $a_n = 8n^3 + n^2 - 2$ show div.

$$\lim_{n \rightarrow \infty} a_n = \infty \therefore \text{seq div}$$

WTS $\forall M > 0, \exists N > 0$ s.t. $\forall n \in \mathbb{N}, n > N \Rightarrow a_n > M$

Let $M > 0$ be arbitrary

$$\text{choose } N = \left(\frac{M}{6}\right)^{\frac{1}{3}} > 0$$

Suppose $n > N$,

$$\begin{aligned} \text{Then } a_n &= 8n^3 + n^2 - 2 \geq 8n^3 - 2 \quad \text{b/c } n^2 > 0 \\ &\geq 8n^3 - 2n^3 \quad \text{b/c } n^3 \geq 1 \\ &= 6n^3 > 6N^3 \\ &= 6 \cdot \left(\left(\frac{M}{6}\right)^{\frac{1}{3}}\right)^3 \\ &= M, \text{ as wanted} \end{aligned}$$

e.g of ②

Prove the seq given $a_1 = \sqrt{6}$ and $a_{n+1} = \sqrt{6 + a_n}$. If $n \geq 1$, then $\{a_n\}$ converges.
(Use BNC1 since $\{a_n\}$ is recursively defined)

WTS $\{a_n\}$ is bounded above and increasing

① WTS $\{a_n\}$ is bounded above

WTS $\exists M \in \mathbb{R}$ s.t. $a_n \leq M \quad \forall n \in \mathbb{N}$

Investigate: $a_1 = \sqrt{6} < \sqrt{9} = 3$ we know $0 < 6 < 9 \Rightarrow \sqrt{6} < \sqrt{9}$

$$a_2 = \sqrt{6 + a_1} = \sqrt{6 + \sqrt{6}} < \sqrt{6 + 3} = 3$$

$$a_3 = \sqrt{6 + a_2} = \sqrt{6 + \sqrt{6 + \sqrt{6}}} < \sqrt{6 + 3} = 3$$

choose $M = 3 \in \mathbb{R}$

WTS $a_n \leq 3 \quad \forall n \in \mathbb{N}$

Base Case: $a_1 = \sqrt{6}$ by def of $\{a_n\}$ $a_1 = \sqrt{6} < \sqrt{9} = 3$ b/c $0 < 6 < 9$ so $\sqrt{6} < \sqrt{9}$

Induction step: WTS $\forall k \in \mathbb{N}$ ($P(k) \Rightarrow P(k+1)$)
holds holds

let $k \in \mathbb{N}$ be arbitrary, assume $a_k \leq 3$ (induction hypothesis)

WTS $a_{k+1} \leq 3$

Consider $a_{k+1} = \sqrt{6+a_k}$ by def of $\{a_n\}$
 ≤ 3 by IH

$$\therefore a_{k+1} \leq \sqrt{6+3} = \sqrt{9} = 3, \text{ as wanted}$$

$\therefore$ By PMI (principle of mathematical induction), $\{a_n\}$ is bounded above by 3 ■

② WTS $\{a_n\}$ is increasing $\Leftrightarrow$ WTS $a_n \leq a_{n+1} \forall n \in \mathbb{N}$

Use derivative analysis for not-recursive fns

Let $n \in \mathbb{N}$ be arbitrary

$$\begin{aligned} \text{Consider } a_n^2 - a_{n+1}^2 &= a_n^2 - (\sqrt{6+a_n})^2, \text{ by def of } \{a_n\} \\ &= a_n^2 - (6+a_n) \\ &= a_n^2 - a_n - 6 \\ &= \underbrace{(a_n-3)}_{\leq 3} \underbrace{(a_n+2)}_{>0} \\ &\leq 0 \end{aligned}$$

$$\therefore a_n^2 - a_{n+1}^2 \leq 0 \quad \forall n \in \mathbb{N}$$

$$\Rightarrow a_n^2 \leq a_{n+1}^2$$

$$\Rightarrow a_n \leq a_{n+1}, \text{ as wanted}$$

$$\therefore \{a_n\} \text{ is } \uparrow.$$

$\therefore$ By BMCT, $\{a_n\}$ converges ■

Proof of Conv property

Let $a_n \rightarrow A$ ① and $b_n \rightarrow B$ ②

(i) $a_n + b_n \rightarrow A+B$

WTS $\forall \epsilon > 0 \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$ then $|a_n + b_n - A - B| < \epsilon$

Let ϵ be arbitrary, choose $N = \max\{N_1, N_2\} > 0$

① $\rightarrow \exists N_1 > 0$ s.t. $\forall n > N_1$, then $|a_n - A| < \frac{\epsilon}{2}$

② $\rightarrow \exists N_2 > 0$ s.t. $\forall n > N_2$, then $|b_n - B| < \frac{\epsilon}{2}$

Suppose $n > N$, then

$$|a_n + b_n - A - B| = |(a_n - A) + (b_n - B)| \leq |a_n - A| + |b_n - B| < \epsilon$$

(ii) $ka_n \rightarrow kA$

WTS $\forall \epsilon > 0 \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$ then $|ka_n - kA| < \epsilon$

Let ϵ be arbitrary. Suppose $n > N$

① $\rightarrow \exists N_1 > 0$ s.t. $\forall n > N_1$, then $|a_n - A| < \frac{\epsilon}{|k|}$

$$|ka_n - kA| = |k| |a_n - A| < |k| \cdot \frac{\epsilon}{|k|} = \epsilon$$

(iii) $a_n b_n \rightarrow AB$

WTS $\forall \epsilon > 0 \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$, then $|a_n b_n - AB| < \epsilon$

Let ϵ be arbitrary, choose $N = \max\{N_1, N_2\} > 0$

Since $b_n \rightarrow B$ then $\exists K \in \mathbb{R}^+$ s.t. $|b_n| < K$

① $\rightarrow \exists N_1 > 0$ s.t. $\forall n > N_1$, then $|a_n - A| < \frac{\epsilon}{2K}$

② $\rightarrow \exists N_2 > 0$ s.t. $\forall n > N_2$, then $|b_n - B| < \frac{\epsilon}{2A}$

$$|a_n b_n - AB| = |b_n(a_n - A) + A(b_n - B)| \leq |b_n| |a_n - A| + |A| |b_n - B| < K |a_n - A| + |A| |b_n - B| < \epsilon$$

$$(iv) \frac{a_n}{b_n} = \frac{A}{B}$$

WTS $\forall \epsilon > 0 \exists N > 0$ s.t. $\forall n \in \mathbb{N}$, if $n > N$, then $|\frac{a_n}{b_n} - \frac{A}{B}| < \epsilon$

Let ϵ be arbitrary. choose $N = \max\{N_1, N_2\} > 0$

$$① \rightarrow \exists N_1 > 0 \text{ s.t. } \forall n > N_1, \text{ then } |a_n - A| < \frac{\epsilon |B|}{4}$$

$$② \rightarrow \exists N_2 > 0 \text{ s.t. } \forall n > N_2, \text{ then } |b_n - B| < \frac{\epsilon B^2}{4|A|}$$

$$|\frac{a_n}{b_n} - \frac{A}{B}| = |\frac{a_n}{b_n} - \frac{A}{b_n} + \frac{A}{b_n} - \frac{A}{B}| = |\frac{a_n - A}{b_n} + \frac{AB - Ab_n}{b_n B}| = |\frac{a_n - A}{b_n} + (B - b_n) \frac{A}{b_n B}| \leq |\frac{1}{b_n}| |a_n - A| + |\frac{A}{b_n B}| |B - b_n|$$

Since $b_n \rightarrow B$, then at some point $|b_n| \geq \frac{|B|}{2} \Rightarrow \frac{1}{|b_n|} \leq \frac{2}{|B|}$

$$\therefore |\frac{1}{b_n}| |a_n - A| + |\frac{A}{b_n B}| |B - b_n| = |\frac{1}{b_n}| |a_n - A| + \frac{2|A|}{|B|^2} |B - b_n| < \frac{2}{|B|} |a_n - A| + \frac{2|A|}{|B|^2} |B - b_n| = \epsilon$$

Proof of if $\{a_n\}$ conv, then its limit is unique

Proof: Suppose $\{a_n\}$ conv to both l_1, l_2 where $l_1, l_2 \in \mathbb{R}$

WTS $\forall \epsilon > 0, |l_1 - l_2| < \epsilon$

$$① \Rightarrow \exists N_1 > 0, \text{ if } n > N_1, \text{ then } |a_n - l_1| < \frac{\epsilon}{2}$$

$$② \Rightarrow \exists N_2 > 0, \text{ if } n > N_2, \text{ then } |a_n - l_2| < \frac{\epsilon}{2}$$

choose $N = \{N_1, N_2\}$, Suppose $n > N$

we have $|l_1 - l_2| = |l_1 - l_2 + a_n - a_n| = |-(a_n - l_1) + (a_n - l_2)| \leq |-(a_n - l_1)| + |a_n - l_2|$ by Δ inequality

$$= |a_n - l_1| + |a_n - l_2|$$

$$< \frac{\epsilon}{2} + \frac{\epsilon}{2}$$

$$= \epsilon, \blacksquare$$

Proof of BMCT

Suppose $\{a_n\}$ is bounded above ①

& $\{a_n\}$ is increasing ② WTS $\{a_n\}$ Conv

WTS $\exists L \in \mathbb{R}, \forall \epsilon > 0 \exists N > 0$ st. $\forall n \in \mathbb{N}$, if $n > N$, then $|a_n - L| < \epsilon$

Define $A = \{a_n | n \in \mathbb{N}\} \subset \mathbb{R}$ and A is not empty since $a_1 \in A$
and A is bounded above by ①

$\therefore \sup(A)$ exist

choose $L = \sup(A) \in \mathbb{R}$ let $\epsilon > 0$ be arbitrary

choose $N \in \mathbb{N}$ and $L - \epsilon < a_n$, suppose $n > N$

we have $a_n \leq L < L + \epsilon$ b/c $\epsilon > 0$

$$\text{and } L - \epsilon < a_n < a_n \leq L < L + \epsilon$$

by ②

$$\therefore L - \epsilon < a_n < L + \epsilon$$

$$- \epsilon < a_n - L < \epsilon$$

$$\therefore |a_n - L| < \epsilon, \blacksquare$$

Series

Ways to determine if a series conv or div

① Inspection and telescoping sum.

② GS Test

③ Div Test

④ Integral Test (IT)

⑤ P-Series Test

① Inspection and telescoping sum

(e.g) a) $\sum_{n=1}^{\infty} \ln\left(\frac{n+1}{n}\right)$ conv or div?

$$\ln\left(\frac{n+1}{n}\right) = \ln(n+1) - \ln(n) \quad S_n = \ln(2) - \ln(1) + \ln(3) - \ln(2) + \ln(4) - \ln(3) + \dots + \ln(n+1) - \ln(n)$$

$$= \ln(n+1) - \ln(1)$$

$$\therefore \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \ln(n+1) = \infty \quad \therefore \text{div}$$

b) $\sum_{n=1}^{\infty} \frac{3}{n^2+3n}$

$$S_n = \frac{3}{n^2+3n} = \frac{3}{n(n+3)} \quad \text{Pfd: } S_n = \frac{3}{n(n+3)} = \frac{A}{n} + \frac{B}{n+3} = \frac{A(n+3)+Bn}{n(n+3)} \quad \therefore A=1 \quad B=-1$$

$$\therefore S_n = \frac{3}{n(n+3)} = \frac{1}{n} - \frac{1}{n+3} \quad \therefore \sum_{n=1}^n S_n = 1 - \frac{1}{4} + \frac{1}{2} - \frac{1}{5} + \frac{1}{3} - \frac{1}{6} + \frac{1}{4} - \frac{1}{7} + \dots + \frac{1}{n-4} - \frac{1}{n-1} + \frac{1}{n-3} - \frac{1}{n} + \frac{1}{n-2} - \frac{1}{n+1} + \frac{1}{n-1} - \frac{1}{n+2} + \frac{1}{n} - \frac{1}{n+3}$$

$$= 1 + \frac{1}{2} + \frac{1}{3} - \frac{1}{n+1} - \frac{1}{n+2} - \frac{1}{n+3}$$

$$\therefore \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} - \frac{1}{n+1} - \frac{1}{n+2} - \frac{1}{n+3}\right)$$

$$= 1 + \frac{1}{2} + \frac{1}{3} = \frac{3}{2} + \frac{1}{3} = \frac{9+2}{6} = \frac{11}{6} \quad \therefore \text{Conv.}$$

② GS Test $\sum_{n=0}^{\infty} ar^n$

For a GS $\sum_{n=1}^{\infty} ar^n$, if $|r| < 1$, then conv to sum $\frac{a}{1-r}$
if $|r| \geq 1$, then div

(e.g)

a) $\sum_{n=2}^{\infty} \frac{\pi(\sqrt{2})^n}{3^{n+1}}$ aside: $3^{n+1} = 3^n \div 3 = 3^n \times \frac{1}{3}$

$$S_n = \frac{\pi(\sqrt{2})^n}{3^{n+1}} = \frac{\pi(\sqrt{2})^n}{3^n \times \frac{1}{3}} = 3\pi \left(\frac{\sqrt{2}}{3}\right)^n$$

$$\therefore r = \frac{\sqrt{2}}{3} \quad \text{and } |r| < 1$$

$$\therefore \text{conv} = \frac{a}{1-r}$$

③ Div Test

if $\lim_{n \rightarrow \infty} a_n \neq 0$, then div

④ IT. Given $\sum a_n$, if $a_n = f(n) \forall n \in \mathbb{N}$ and $f(x)$ is cont, positive, decreasing on $[1, \infty)$ then $\sum_{n=1}^{\infty} a_n \Leftrightarrow \int_1^{\infty} f(x) dx$
div $\Leftrightarrow$ div
conv $\Leftrightarrow$ conv

(eg) $\sum_{n=0}^{\infty} \frac{1}{1+4n^2}$ conv or div?

Let $f(n) = \frac{1}{1+4n^2}$, then $f(x) = \frac{1}{1+4x^2}$. (consider $f(x)$ on $[1, \infty)$)

• $f(x) > 0$ and $f'(x) \leq 0$

$\therefore f(x)$ is positive, $\downarrow$ and cont on $[1, \infty)$

consider $\int_0^{\infty} f(x) dx = \lim_{A \rightarrow \infty} \int_0^A \frac{1}{1+4x^2} dx = \lim_{A \rightarrow \infty} \left. \frac{\arctan(2x)}{2} \right|_0^A = \frac{\pi}{4}$ $\therefore$ limit exist $\therefore \int_0^{\infty} f(x) dx$ conv
by IT, series conv.

⑤ p-series Test

Given $\sum_{n=1}^{\infty} \frac{1}{n^p}$. if $p > 1$ then conv
if $0 < p \leq 1$ then div

⑥ CT for series

if $0 \leq a_n \leq b_n \forall n \in \mathbb{N}$, b_n conv $\Rightarrow a_n$ conv

if $0 \leq c_n \leq a_n \forall n \in \mathbb{N}$, c_n conv $\Rightarrow a_n$ conv

⑦ LCT* Let $\sum a_n$ $\sum b_n$ be series. Suppose $a_n b_n > 0 \forall n \in \mathbb{N}$

Defined $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = p$

① if $0 < p < \infty \Rightarrow$ both conv or div

② if $p = 0$ and $\sum b_n$ conv $\Rightarrow \sum a_n$ conv

③ if $p = \infty$ and $\sum b_n$ div $\Rightarrow \sum a_n$ div

⑧ Alternating Series Test (AST)

Given $\sum (-1)^n b_n$, $b_n > 0$

if ① $b_n \geq b_{n+1} > 0 \forall n \in \mathbb{N}$

② $\lim_{n \rightarrow \infty} b_n = 0$ (check this first)

then conv

e.g

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{ cc? Ac? div?}$$

$$\text{consider } \sum_{n=1}^{\infty} \left| \frac{(-1)^n}{n} \right| = \sum_{n=1}^{\infty} \frac{1}{n} \quad p=1 \Rightarrow \text{div}$$

$$\text{consider } \sum_{n=1}^{\infty} \frac{(-1)^n}{n} = \sum_{n=1}^{\infty} (-1)^n \cdot \frac{1}{n}$$

$$\textcircled{1} \lim_{n \rightarrow \infty} \frac{1}{n} = 0$$

② let $n \in \mathbb{N}$ be arbitrary

we know $0 < n < n+1$

$$\Rightarrow \frac{1}{n} > \frac{1}{n+1}$$

$\therefore$ conv

$\therefore$ cc

⑨ Ratio Test

Let $\sum a_n$ be a series with $a_n \in \mathbb{R} - \{0\}$

$$\text{Define } p = \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} \quad \text{if } p < 1 \Rightarrow \sum a_n \text{ Ac and conv}$$

$$p > 1 \Rightarrow \sum a_n \text{ div}$$

$$p = 1 \Rightarrow \text{we don't know}$$

⑩ Power Series

A series in the form of $\sum_{n=0}^{\infty} C_n (x-a)^n$

eg. for what $x \in \mathbb{R}$ does $\sum_{n=1}^{\infty} \frac{(-1)^n (x-2)^n}{n 4^n}$ conv?

(i.e) what is the interval of conv

$$I = \{x \in \mathbb{R}, \sum_{n=1}^{\infty} \frac{(-1)^n (x-2)^n}{n 4^n} \text{ conv}\}$$

① RT

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{C_{n+1} (x-2)^{n+1}}{C_n (x-2)^n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} (x-2)^{n+1}}{(n+1) 4^{n+1}} \cdot \frac{n 4^n}{(-1)^n (x-2)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^n (-1)}{(-1)^n} \cdot \frac{4^n}{4^n \cdot 4} \cdot \frac{n}{n+1} \cdot \frac{(x-2)^{n+1}}{(x-2)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| (-1) \cdot \frac{1}{4} \cdot \frac{n}{n+1} \cdot (x-2) \right| \\ &= \frac{1}{4} |x-2| \lim_{n \rightarrow \infty} \frac{n}{n+1} \\ &= \frac{|x-2|}{4} \\ \therefore \text{by RT } \frac{|x-2|}{4} < 1 &\Rightarrow \text{conv} \\ |x-2| < 4 &\text{ radius} \end{aligned}$$

② check end pts $x=a+r, x=a-r$

a) $x=a+r=2+4=6$

$$\sum_{n=1}^{\infty} \frac{(-1)^n (4)^n}{n 4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \quad \checkmark$$

b) $x=a-r=2-4=-2$

$$\sum_{n=1}^{\infty} \frac{(-1)^n (-4)^n}{n 4^n} = \sum_{n=1}^{\infty} \frac{1}{n}$$

Series Test summary

- ① Inspection and telescoping ($\ln(\frac{n+1}{n}) = \ln(n+1) - \ln(n)$) *conv to # or div*
- ② GS Test $\sum_{n=1}^{\infty} ar^n$. if $|r| < 1 \Rightarrow \text{conv to } \frac{a}{1-r}$ a is the first nonzero term in GS *conv to # or div*
if $|r| \geq 1 \Rightarrow \text{div}$

③ Div test

if $\lim_{n \rightarrow \infty} a_n \neq 0 \Rightarrow \text{div}$ *div*

④ Integral Test

Given $\sum a_n, a_n = f(n)$. if $f(x)$ cont., +, $\downarrow$ on $[1, \infty)$

Then $\int_1^{\infty} f(x) dx \text{ conv} \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=1}^n a_k \text{ conv}$ *conv or div*

$\int_1^{\infty} f(x) dx \text{ div} \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=1}^n a_k \text{ div}$

⑤ p-Series Test

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ if $p > 1 \Rightarrow \text{conv}$ *conv or div*
if $0 < p \leq 1 \Rightarrow \text{div}$

⑥ CT $\sum a_n, \sum b_n, \sum c_n$

if $0 \leq a_n \leq b_n \forall n \in \mathbb{N}$ and $\sum b_n \text{ conv} \Rightarrow \sum a_n \text{ conv}$ *conv or div*

if $0 \leq c_n \leq a_n \forall n \in \mathbb{N}$ and $\sum c_n \text{ div} \Rightarrow \sum a_n \text{ div}$

⑦* LCT Suppose $a_n b_n > 0 \forall n \in \mathbb{N}$

Defined $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = p$

① if $0 < p < \infty \Rightarrow \text{both conv or div}$ *conv or div*

② if $p = 0$ and $\sum b_n \text{ conv} \Rightarrow \sum a_n \text{ conv}$

③ if $p = \infty$ and $\sum b_n \text{ div} \Rightarrow \sum a_n \text{ div}$

⑧ Alternating Series Test (AST)

Given $\sum (-1)^n b_n, b_n > 0$

if ① $b_n \geq b_{n+1} > 0 \forall n \in \mathbb{N}$

② $\lim_{n \rightarrow \infty} b_n = 0$ (check this first)

then conv

$$P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$

⑨ Ratio Test

Let $\sum a_n$ be a series with $a_n \in \mathbb{R} - \{0\}$

Define $\rho = \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}$ if $\rho < 1 \Rightarrow \sum a_n \text{ AC and conv}$

$\rho > 1 \Rightarrow \sum a_n \text{ div}$

$\rho = 1 \Rightarrow \text{we don't know}$

if $\rho = 0 \Rightarrow \text{infinite radius}$
and interval is $\mathbb{R}$

if $\rho = \infty \Rightarrow \text{radius is } 0$
and $I = \{a\}$